

10050 Auburn Park Drive Fort Wayne, IN 46825 260-4-FAMILY

Patient Name: ______

Date: _____ Today's Weight: _____

We recommend giving a full dose of Acetaminophen with a full dose of Ibuprofen at the same time every 6 hours.

Acetaminophen (Tylenol/Genexa)								
Child's Weight (Pounds)	6-11	12-17	18-23	24-35	36-47	48-59	60-71	lbs
Syrup: 160mg/5 milliliters (mL)	1.25	2.5	3.75	5	7.5	10	12.5	mL
Chewable: 80-mg tablets			1 ½	2	3	4	5	tablets
Chewable: 160-mg tablets				1	1 ½	2	2 ½	tablets

Frequency: Repeat every 4-6 hours as needed.

Caution: Don't give more than 5 doses in 24-hour period.

Ibuprofen (Advil, Motrin)									
Child's Weight (Pounds)	12-17	18-23	24-35	36-47	48-59	60-71	lbs		
Infant Drops: 50mg/1.25mL	1.25	1.875	2.5	3.75	5		mL		
Liquid: 100mg/5 milliliters (mL)	2.5	4	5	7.5	10	12.5	mL		
Chewable: 50 -mg tablets			2	3	4	5	tablets		
Jr. Strength: 100-mg tablets					2	2 ½	tablets		

Frequency: Repeat every 6-8 hours as needed.

Caution: Don't use under 6 months of age.

Diphenhydramine (Benadryl)								
Child's Weight (pounds)	20-24	25-37	38-49	50-99	100+	lbs		
Liquid: 12.5mg/ 5 milliliters (mL)	4	5	7.5	10		mL		
Chewable: 12.5 mg		1	1 ½	2	4	tablets		
Tablets: 25 mg		1/2	1/2	1	2	tablets		

Frequency: Repeat every 6-8 hours as needed.

Caution: Don't use under 6 months of age.